

The Effect of Condensates on the Characterization of Transiting Planet Atmospheres with Transmission Spectroscopy

Jonathan J. Fortney^{*}

Space Science and Astrobiology Division, NASA Ames Research Center, MS 245-3, Moffett Field, CA 94035

Accepted September 7, 2005

ABSTRACT

Through a simple physical argument we show that the slant optical depth through the atmosphere of a ‘hot Jupiter’ planet is ~ 35 –90 times greater than the normal optical depth. This not unexpected result has direct consequences for the method of transmission spectroscopy for characterizing the atmospheres of transiting giant planets. The atmospheres of these planets likely contain minor condensates and hazes which at normal viewing geometry have negligible optical depth, but at slant viewing geometry have appreciable optical depth that can obscure absorption features of gaseous atmospheric species. We identify several possible condensates. We predict that this is a general masking mechanism for all planets, not just for HD 209458b, and will lead to weaker than expected or undetected absorption features. Constraints on an atmosphere from transmission spectroscopy are not the same as constraints on an atmosphere at normal viewing geometry.

Key words: planetary systems; radiative transfer.

1 INTRODUCTION

To date, 8 extrasolar giant planets (EGPs) are known to transit their parent stars in tight, several day orbits. Characterizing the atmospheres of two of these planets (HD 209458b and TrES-1) has been a major goal for many astronomers in the past few years. Not long after the discovery of the transits of planet HD 209458b (Charbonneau et al. 2000; Henry et al. 2000), a number of studies appeared in the literature on radiative transfer aspects of ‘Pegasi planet’ (or ‘hot Jupiter’) transits, namely how stellar light passing through a planet’s atmosphere can be absorbed at wavelengths where opacity is high. This would lead a distant observer who obtained a spectrum of the star during a transit of the planet to see the planet’s atmospheric absorption spectrum superimposed on the star’s spectrum. The physics behind this process were laid out theoretically and modeled by Seager & Sasselov (2000), Brown (2001), and Hubbard et al. (2001). The consensus of these three studies was that for HD 209458b, for a clear, cloudless atmosphere, that the transit depth (0.016 in relative flux) could itself vary by up to a few % due to absorption by gaseous sodium, potassium, water, and carbon monoxide.

Shortly thereafter Charbonneau et al. (2002) used the

STIS instrument aboard *Hubble Space Telescope* to observe the predicted sodium absorption doublet at 589 nm. However, the magnitude of this absorption was 2–3 times weaker than had been predicted by theoretical models that assumed a clear and cloudless atmosphere. A number of possible reasons for this discrepancy were put forth in Charbonneau et al. (2002), including a global underabundance of sodium, ionization of sodium by stellar flux, sodium being tied up in condensates or molecules (see also Atreya et al. 2003), or high clouds obscuring the absorption of sodium (see also Seager & Sasselov 2000). In addition, Barman et al. (2002) found that neutral atomic sodium may be out of local thermodynamic equilibrium, leading to a deficit of sodium atoms able to absorb at 589 nm, relative to LTE calculations. Fortney et al. (2003) derived a self-consistent pressure-temperature (P – T) profile for HD 209458b, and found that silicate and iron clouds reside high in the planet’s atmosphere, at the several mbar level, and that these opaque clouds mask the absorption of sodium enough to match the Charbonneau et al. (2002) observations. However, as noted by the authors at the time, this conclusion is extremely sensitive to the P – T profile calculated for the atmosphere. Recently Iro et al. (2005) have shown that a substantial day-night temperature contrast in the planet’s atmosphere could lead to a sink of atomic

^{*} E-mail: jfortney@arc.nasa.gov

sodium, as the atom could be tied up into the condensate Na_2S on the planet's night side.

In addition to the sodium observation, Deming et al. (2005) have recently attempted to observe absorption due to first overtone bands of CO at $\sim 2.3 \mu\text{m}$, using NIRSPEC on the Keck Telescope. Their sensitivity was high enough such that if CO was present in the abundances predicted for a clear and cloud-free atmosphere, that the CO should have been detected. However, it was not, and the authors point to the masking effect of high clouds as the likely culprit. Currently other searches are underway to detect the absorption of H_2O (Harrington et al. 2002) and H_3^+ (Haywood et al. 2004) in the atmosphere of HD 209458b. Narita et al. (2005) have also reported upper limits due to absorption by Li, H, Fe, and Ca. These studies are in addition to the exosphere absorption studies such as Vidal-Madjar et al. (2003, 2004), which have yielded important results, but are not the subject of our study here.

To date, there has been no discussion in the extrasolar planets literature regarding how condensates less abundant than the 'standard' brown dwarf condensates (such as silicates and iron) may effect transmission spectroscopy. In the following sections we show that condensates that may have insignificant optical depth when viewed at normal geometry can have appreciable optical depths for the *slant* viewing geometry relevant for transits. Searches for transmission absorption features in the atmosphere of HD 209458b, or any other similar searches for absorption during transits of other planets in the future, will very likely find weak or non-existent absorption features. Constraints on atmospheric abundances derived from transmission spectroscopy will not map directly as constraints on abundances under normal viewing geometry.

2 GEOMETRY OF THE PROBLEM

The studies of Hubbard et al. (2001) and Fortney et al. (2003) showed that the pressures probed by transit observations are sensitive function of wavelength. Specifically for HD 209458b, Fortney et al. (2003) showed that in the spectral region from 580 to 640 nm, which spanned the Charbonneau et al. (2002) observations, the pressure where the slant optical depth reached 1 varied from a few μbar near the sodium line cores, to potentially 50 mbar at 640 nm, if the atmosphere was clear. This pressure could reach \sim a few hundred mbar at wavelengths of relatively low opacity.

At these low pressures it is reasonable, to first order, to approximate the atmosphere as having a constant temperature with height. If a planet's atmosphere is in hydrostatic equilibrium, with a constant temperature and mean molecular weight, then the barotropic law holds, which states (Chamberlain & Hunten 1987):

$$p(r) = p(r_0) \exp\left(-\frac{GMm}{kTr_0}(r - r_0)\right) \quad (1)$$

where p is the pressure, r is the radius of interest, r_0 is the reference radius, G is the gravitational constant, M is the mass of the planet, m is the mean mass of a molecule in the atmosphere, k is Boltzmann's constant, and T is the temperature. The vertical integrated column density of the

atmosphere $N_V(r_0)$, above a given local density $n(r_0)$, is given by:

$$N_V(r_0) \equiv \int_{r_0}^{\infty} n(r) dr = \int_0^{p(r_0)} \frac{r^2}{GMm} dp \approx \frac{p(r_0)}{g(r_0)m} = n(r_0)H, \quad (2)$$

where H is the scale height ($H = kT/mg$) and g is the planet's gravitational acceleration. Clearly, if our reference density is n_0 , our column density is then n_0H . With the assumption that g is constant in the atmosphere Eq. (1) can be simplified to:

$$p(z) \approx p(z_0) \exp\left(-\frac{z - z_0}{H}\right) \quad (3)$$

where here we have replaced the two radii, r and r_0 with heights from a given level, z and z_0 , respectively.

We now turn to Fig. 1. Here a is a given radius of the planet, say where the normal optical depth is unity. The thickness of our atmosphere is z and x is a line tangential to our optical depth unity 'surface,' horizontal to the horizon. Using the Pythagorean theorem and that $2az \gg z^2$, then $z \approx x^2/2a$. Therefore, Eq. (3), when written in terms of density, rather than pressure, becomes

$$n(x) = n_0 \exp\left(\frac{-x^2}{2aH}\right) \quad (4)$$

and the horizontal integrated density N_H , from horizon to horizon, is

$$N_H = \int_{-\infty}^{\infty} n(x) dx = n_0 \sqrt{2\pi aH} \quad (5)$$

The ratio of the horizontal integrated density to the vertical integrated density is the quantity of interest here. This quantity, which we will label η , simplifies to

$$\eta = \frac{N_H}{N_V} = \frac{\tau_H}{\tau_V} = \sqrt{\frac{2\pi a}{H}}. \quad (6)$$

This shows that the horizontal integrated density is significantly larger than the vertical integrated density. This ratio is ~ 75 for Earth (and ~ 128 for Jupiter), and leads to an atmosphere that has significantly different absorption and scattering properties at a *slant* geometry than for *normal* geometry, which one can easily observe at sunset. Since the optical depth is directly proportional to the column density, η is also the ratio of the slant optical depth to the normal optical depth.

3 APPLICATION TO EGPS

3.1 Atmospheric Properties

For HD 209458b, assuming $T = 1200 \text{ K}$, $a = 10^5 \text{ km}$, and mean molecular weight $\mu = 2.3$, we find $\eta = 38$. For TrES-1, which although similar in mass has a 30% smaller radius and is $\sim 300 \text{ K}$ colder in effective temperature (Fortney et al. 2005), we find $\eta = 50$. A minor condensate, having a normal optical depth of 0.02, which would be easily ignored when calculating a planet's emission or reflection spectrum, would have an optical depth of 1 in slant transmission through the planet's atmosphere.

In Fig. 2 we plot self-consistent P - T profiles for planets HD 209458b and TrES-1, as taken from Fortney et al.




Figure 1. Diagram of slant vs. normal geometry. a is the planet’s radius out to a standard level, say, the radius at normal optical depth unity. z is the thickness of the atmosphere above this level, out to some very low pressure. x is the distance to this same low pressure, towards the horizon.

(2005). Also plotted are condensation curves for a variety of equilibrium condensates, spanning a large range in temperature. The condensation curves are taken from Lodders & Fegley (2005). We note that the P - T profile shown here for HD 209458b is cooler than that of Fortney et al. (2003) because here we assume the planet is able to reradiate absorbed stellar flux over the entire planet, whereas Fortney et al. (2003) assumed this reradiation could only occur on the planet’s day side, leading to a warmer day side P - T profile. In addition, Fortney et al. (2003) utilized a different model atmosphere program (see Sudarsky et al. 2003).

We have used the transit radiative transfer program described in Hubbard et al. (2001) and Fortney et al. (2003) to compute both the normal optical depth and slant optical depth at various pressures in the atmosphere of both planets. This code takes a computed model atmosphere, for which T , P , ρ , and extinction cross-sections are known as a function of altitude, under the assumption of hydrostatic equilibrium, and places the atmosphere onto an opaque disk of a given radius. Normal and slant optical depths are then computed numerically through the atmosphere. For isothermal atmospheres, we recover the same ratio η we calculated analytically, within less than $\sim 1\%$. For the atmospheres we consider here our pressure range of interest is the upper troposphere, and temperature decreases with altitude. Integrating upwards in altitude from a pressure of 1 bar, we calculate $\eta = 30$ for HD 209458b and 40 for TrES-1, about 80% of our simple analytic case. However, integrating from a lower pressure, say 10 mbar, may be more relevant, and from this pressure we calculate $\eta = 35.5$ for HD 209458b and 47 for TrES-1, which is $\sim 94\%$ of the value obtained from our simple analytical treatment. In summary, a more detailed analysis gives ratios of the slant optical depth to normal optical depth for these atmospheres that is consistent with our earlier simple analysis. These values are collected in Table 1.

3.2 Condensate Scale Height

The values for η calculated so far assume that the scale height for condensate is the same as the scale height of the surrounding gas. However, there is strong evidence in

Jupiter’s atmosphere that its visible ammonia cloud is more compact than the surrounding gas. Using Voyager infrared spectra Carlson et al. (1994) derived a ratio of the scale height of condensate (H_{cond}) to scale height of gas (H) of 0.35 ± 0.10 for Equatorial Zones and 0.40 ± 0.10 for Northern Tropical Zones. Observations of the Jovian tropics with the *Infrared Space Observatory* by Brooke et al. (1998) indicate a scale height ratio of 0.3.

Further evidence for clouds of small vertical extent comes from observations of L-type stars and brown dwarfs, which have silicate and iron clouds in their visible atmospheres. The observed spectra of these objects have been accurately modeled by Marley et al. (2002) and Marley et al. (2004) using the 1D cloud model of Ackerman & Marley (2001) with a sedimentation efficiency parameter, $f_{sed} = 2$ -3. This f_{sed} range gives silicate and iron clouds with $H_{cond}/H = 0.25 - 0.30$ (Ackerman & Marley 2001). These observations and models indicate that equilibrium condensates, across a wide range of temperatures and chemical species, have scale heights that are $\sim 1/3$ of the local gas scale height. This leads to values of η that are $\sim 75\%$ larger than calculated earlier. Values of η would then increase to 66 for HD 209458b and 87 for TrES-1.

3.3 Minor Condensates and Hazes

For brown dwarf atmospheres, corundum (Al_2O_3), iron (Fe), and silicates (MgSiO_3 and/or Mg_2SiO_4) appear to be the only condensates that have appreciable optical depth, and therefore leave any imprint on the spectra of these objects. (See, for instance, Ackerman & Marley 2001; Marley et al. 2002; Cooper et al. 2003; Allard et al. 2001; Lodders & Fegley 2005). However, at slant viewing geometry, one likely has to consider condensates that may be a factor of 100-1000 less abundant, compared to the silicates. From the work of (Fegley & Lodders 1994), on condensation in the deep atmospheres of Jupiter and Saturn, and Lodders (2003), on the condensation temperatures of the elements, we can highlight condensates that fit into this abundance range. In order of decreasing condensation temperature, these are: Cr, MnS, Na_2S , ZnS, KCl, and $\text{NH}_4\text{H}_2\text{PO}_4$. These are the condensation curves plotted in Fig. 2.

To analyse the potential optical depths of these species, we will closely follow the analysis of Marley (2000), who derived an expression for the maximum optical depth of condensates in substellar atmospheres. We will use this equation to determine the relative optical depths of various species. This expression is:

$$\tau_\lambda = 75\epsilon Q_\lambda(r_c)\varphi \left(\frac{P_c}{1 \text{ bar}} \right) \left(\frac{10^5 \text{ cm sec}^{-2}}{g} \right) \left(\frac{1\mu\text{m}}{r_c} \right) \left(\frac{1.0\text{g cm}^{-3}}{\rho_c} \right), \quad (7)$$

where ϵ is a factor ≤ 1 that accounts for the finite amount of species left over after condensation, because of the vapor pressure above the condensate, Q_λ is the wavelength dependent extinction efficiency from Mie theory, r_c is the radius of the condensate particles, and $\varphi = f m_c / \bar{m}$, where f is the mixing ratio of the species, m_c is the molecular weight of the condensed species, \bar{m} is the mean molecular weight of the atmosphere. Additionally, P_c is the condensation pressure, g is gravitational acceleration in the atmosphere, and ρ_c is the mass density of the condensate. Since at this point we are only interested in the *relative* optical depths of the various

species at a given P_c , we will make several simplifications. Following Marley, we assume that ϵ , Q_λ , r_c , ρ_c are approximately equal for the condensate species, then the optical depth ratio for given condensates 1 and 2 reduces to

$$\frac{\tau_1}{\tau_2} = \frac{f_1 m_{c1}}{f_2 m_{c2}}. \quad (8)$$

For illustrative purposes, we used a P - T profile for HD 209458b that is slightly warmer than that shown in Fig. 2, along with the Ackerman & Marley (2001) cloud model with $f_{sed} = 2$, to compute the normal optical depth of a MgSiO_3 cloud with a base at 30 mbar. We find that the optical depth is 0.5 at normal viewing geometry at a wavelength of $1 \mu\text{m}$. Based on this calculated value, we then determine normal optical depths for our other condensates, if they were to form at this pressure, using Eq. 8. We also calculated the slant optical depths for HD 209458b, assuming $\eta = 66$. These are listed in Table 2. The f values for each condensate is the f of the limiting atomic species for each condensate, as given for ‘Solar System Abundances’ in Lodders (2003). This assumes that there are not other molecules tying up the atoms, an assumption that is accurate at these high temperatures. The slant optical depths calculated are on the order of ~ 0.1 -1, meaning their opacity is not negligible. From this analysis we can see that these minor condensates may well mask absorption features due to gaseous atomic and molecular species. In addition, if the atmospheres of transiting planets are enhanced in heavy elements these optical depths could well be larger. Jupiter’s atmosphere is enhanced in heavy elements by a factor of about 3 times over solar composition (Atreya et al. 2003) and Saturn’s methane abundance has recently been pinned at ~ 7 times solar (Flasar et al. 2005).

Non-equilibrium hazes, such as the photochemically-produced hydrocarbon hazes found in the atmospheres of Jupiter, Saturn, Uranus, Neptune, Titan, and Los Angeles, have long been observed and modeled (e.g., for Jupiter: West et al. 1986; Tomasko et al. 1986; Rages et al. 1999). In Jupiter these stratospheric hazes can have normal optical depths of a few tenths at high latitudes. The scale height of these hazes is generally similar to the scale height of the surrounding gas (Moses et al. 1995; Rages et al. 1999).

To date, only Liang et al. (2004) have studied whether photochemically-produced hydrocarbon hazes will be found in Pegasi planet atmospheres. These authors found it very unlikely that Pegasi planets would have hazes of this sort, due to several reasons. These include: a lack of methane to be photolyzed (CO is the dominant carbon carrier), the high atmospheric temperatures, which would not allow any hydrocarbon products that were formed to condense, and fast reverse reactions that quickly break down hydrogenated carbon compounds. However, these authors acknowledge they do not consider ion-neutral chemistry, and they also note that the stellar wind could be a vast source for high-energy charged particles. Much work still needs to be done to definitively rule out hazes not predicted by equilibrium chemistry. Even condensates that form relatively thin hazes could have important effects on transmission spectroscopy.

4 DISCUSSION AND CONCLUSIONS

In our calculations we have used a relatively simple atmosphere model. We assume that the limb of the planet is uniform around the entire planet and that the atmosphere has a uniform P - T profile. It is certainly possible that this condition will not be met in reality (see Showman & Guillot 2002; Cho et al. 2003; Cooper & Showman 2005; Barman et al. 2005). As Iro et al. (2005) showed in the region of the 589 nm sodium doublet, it is possible for ‘hot’ and ‘cold’ limbs of a planet to show significantly different transmission signatures. Our main argument will likely hold even if the distribution of condensates is more complex than we have assumed. For instance, if the temperature on the limb simply monotonically increases from the night to day sides, one could imagine a series of atmosphere profiles where a given condensation curve is crossed at progressively lower pressures. At a given pressure at the terminator, this could lead to a lower condensate slant optical depth on the night side, but a higher slant optical depth on the day side, relative to our treatment here. It is certainly possible that different condensates could be important in different locations in a planet’s atmosphere.

In this paper we have taken a straightforward look at the optical depth that minor condensates may have in the slant viewing geometry relevant to planetary transits. While our findings are potentially not unanticipated, we felt the need to discuss this issue because the impact of minor condensates on the transit characterization of planetary atmospheres had not been discussed in the extrasolar planet literature to date. Our conclusions can be summarized as follows:

- For the *slant* viewing geometry relevant to transmission spectroscopy observations of EGPs, the slant optical depth can be on the order of 35-90 times larger than the normal optical depth. This depends upon the scale height of the condensate specifically, which may be smaller than the scale height of the surrounding gas.
- Constraints on cloud location and thickness, and/or constraints on chemical abundances, obtained from transmission spectroscopy will not map directly onto constraints for the atmosphere when viewed at normal geometry. A cloud can be optically thick at slant viewing geometry and optically thin at normal viewing geometry. Thus, there could be abundant atomic Na and CO in the atmosphere of HD 209458b, even though Charbonneau et al. (2002) observed only a weak Na absorption feature, and Deming et al. (2005) observed no CO. The obscuring opacity source for this planet could be condensed Cr, MnS, silicates, or Fe, and will depend on the actual temperatures on the planet’s limb.
- Minor equilibrium condensates or photochemically derived hazes that may be reasonably ignored for normal viewing geometry due to their low optical depths may have to be taken into account at slant viewing geometry.
- These minor condensates may include ones we could reasonably predict the location and distribution of (such as MnS) with an accurate pressure-temperature profile and assumed chemical mixing ratios, and those that we may remain ignorant of, such as photochemically produced hazes, that may have negligible *normal* optical depths.

We assert that transmission spectroscopy will continue to yield abundances of expected chemical species far below

those predicted for a ‘clear’ atmosphere, for HD 209458b, and for other planets that may be studied in the future.

ACKNOWLEDGMENTS

I thank K. Lodders for numerous helpful discussions and for sharing condensation data prior to publication. I thank M. S. Marley for comments on the manuscript and also D. Charbonneau, T. M. Brown, J. W. Barnes, C. S. Cooper, and W. B. Hubbard for interesting discussions. J. J. F. is supported by an NRC Research Associateship.

REFERENCES

- Ackerman A. S., Marley M. S., 2001, *ApJ*, 556, 872
Allard F., Hauschildt P. H., Alexander D. R., Tamanai A., Schweitzer A., 2001, *ApJ*, 556, 357
Atreya S. K., Mahaffy P. R., Niemann H. B., Wong M. H., Owen T. C., 2003, *Planet. Space Sci.*, 51, 105
Barman T. S., Hauschildt P. H., Allard F., 2005, *ApJ*, in press (astro-ph/0507136)
Barman T. S., Hauschildt P. H., Schweitzer A., Stancil P. C., Baron E., Allard F., 2002, *ApJ*, 569, L51
Brooke T. Y., Knacke R. F., Encrenaz T., Drossart P., Crisp D., Feuchtgruber H., 1998, *Icarus*, 136, 1
Brown T. M., 2001, *ApJ*, 553, 1006
Carlson B. E., Lacis A. A., Rossow W. B., 1994, *JGR*, 99, 14623
Chamberlain J. W., Hunten D. M., 1987, *Orlando FL Academic Press Inc International Geophysics Series*, 36
Charbonneau D., Brown T. M., Latham D. W., Mayor M., 2000, *ApJ*, 529, L45
Charbonneau D., Brown T. M., Noyes R. W., Gilliland R. L., 2002, *ApJ*, 568, 377
Cho J. Y.-K., Menou K., Hansen B. M. S., Seager S., 2003, *ApJ*, 587, L117
Cooper C. S., Showman A. P., 2005, *ApJ*, 629, L45
Cooper C. S., Sudarsky D., Milsom J. A., Lunine J. I., Burrows A., 2003, *ApJ*, 586, 1320
Deming D., Brown T. M., Charbonneau D., Harrington J., Richardson L. J., 2005, *ApJ*, 622, 1149
Fegley B. J., Lodders K., 1994, *Icarus*, 110, 117
Flasar F. M., et al., 2005, *Science*, 307, 1247
Fortney J. J., Marley M. S., Lodders K., Saumon D., Freedman R., 2005, *ApJ*, 627, L69
Fortney J. J., Sudarsky D., Hubeny I., Cooper C. S., Hubbard W. B., Burrows A., Lunine J. I., 2003, *ApJ*, 589, 615
Harrington J., Deming D., Matthews K., Richardson L. J., Rojo P., Steyert D., Wiedemann G., Zeehandelaar D., 2002, *Bulletin of the American Astronomical Society*, 34, 1175
Haywood J., Gibb E., Brittain S., Rettig T., Kulesa C., 2004, *AAS/Division for Planetary Sciences Meeting Abstracts*, 36, #35.02
Henry G. W., Marcy G. W., Butler R. P., Vogt S. S., 2000, *ApJ*, 529, L41
Hubbard W. B., Fortney J. J., Lunine J. I., Burrows A., Sudarsky D., Pinto P., 2001, *ApJ*, 560, 413
Iro N., Bezdard B., Guillot T., 2005, *A&A*, 436, 719

Table 1. Quantities of Interest for Various Planets

Planet	a (km)	H (km)	η	H_{cond} (km)	η_{cond}^a
Earth	6400	7	76	2.3	132
Jupiter	70000	27	128	9	221
HD 209458b	100000	440	38	147	66
TrES-1	75000	185	50	62	87

^a Value for H_{cond} and η_{cond} (condensate) are computed assuming $H_{cond} = 1/3 H_{gas}$.

Table 2. Optical depths (at a wavelength of 1 μm) of lesser condensates in HD 209458b, for an atmosphere of solar composition

Condensate	Molecular Weight	Limiting Species	f	normal τ	Slant τ
MgSiO ₃	100.4	Si/Mg	7×10^{-5}	0.5	33
Na ₂ S	78.1	Na	2×10^{-6}	0.011	0.73
NH ₄ H ₂ PO ₄	113	P	5.8×10^{-7}	0.011	0.73
MnS	87	Mn	6.3×10^{-7}	0.0093	0.61
Cr	85	Cr	8.8×10^{-7}	0.0077	0.51
KCl	74.6	K	2.5×10^{-7}	0.0028	0.18
ZnS	97.5	Zn	8.5×10^{-8}	0.0014	0.09

- Liang M., Seager S., Parkinson C. D., Lee A. Y.-T., Yung Y. L., 2004, *ApJ*, 605, L61
Lodders K., 2003, *ApJ*, 591, 1220
Lodders K., Fegley B., 2005, *Astrophysics Update*, in press
Marley M. S., 2000, in *ASP Conf. Ser. 212: From Giant Planets to Cool Stars*, Ed. C. A. Griffith and M. S. Marley, p. 152
Marley M. S., Cushing M. C., Saumon D., 2004, 13th Workshop on Cool Stars, Stellar Systems, and the Sun, eds. F. Favata, G. A. J. Hussain, & B. Battrock, ESA SP-560, p. 794
Marley M. S., Seager S., Saumon D., Lodders K., Ackerman A. S., Freedman R. S., Fan X., 2002, *ApJ*, 568, 335
Moses J. I., Rages K., Pollack J. B., 1995, *Icarus*, 113, 232
Narita N., Suto Y., Winn J. N., Turner E. L., Aoki W., Leigh C. J., Sato B., Tamura M., Yamada T., 2005, *PASJ*, 57, 471
Rages K., Beebe R., Senske D., 1999, *Icarus*, 139, 211
Seager S., Sasselov D. D., 2000, *ApJ*, 537, 916
Showman A. P., Guillot T., 2002, *A&A*, 385, 166
Sudarsky D., Burrows A., Hubeny I., 2003, *ApJ*, 588, 1121
Tomasko M. G., Karkoschka E., Martinek S., 1986, *Icarus*, 65, 218
Vidal-Madjar A., Désert J.-M., Lecavelier des Etangs A., Hébrard G., Ballester G. E., Ehrenreich D., Ferlet R., McConnell J. C., Mayor M., Parkinson C. D., 2004, *ApJ*, 604, L69
Vidal-Madjar A., Lecavelier des Etangs A., Désert J.-M., Ballester G. E., Ferlet R., Hébrard G., Mayor M., 2003, *Nature*, 422, 143
West R. A., Strobel D. F., Tomasko M. G., 1986, *Icarus*, 65, 161




Figure 2. Pressure-temperature profiles for TrES-1 and HD 209458b. Condensation curves for various compounds, as taken from Lodders & Fegley (2005) are shown as dotted lines. The boundary where CH_4 and CO have the same abundance is shown as a dashed line.

This paper has been typeset from a $\text{T}_\text{E}\text{X}$ / $\text{L}^\text{A}\text{T}_\text{E}\text{X}$ file prepared by the author.